



ALOPECIA (HAIR LOSS)

Alopecia can be caused by internal or external factors. Although there are various types of alopecia there are three main groups which are: AREATE, DIFUSE AND ANDROGENETIC

Dental focus							
Upper Jaw	Incisors frontal Canines frontal and parietals Pre-molars frontal. parietal and temporal						
Lower Jaw	<table border="0" style="width: 100%;"> <tr> <td>Incisors (men)</td> <td style="text-align: right;">chin</td> </tr> <tr> <td>Canines (men)</td> <td style="text-align: right;">chin</td> </tr> <tr> <td>Pre-molars (men)</td> <td style="text-align: right;">chin</td> </tr> </table>	Incisors (men)	chin	Canines (men)	chin	Pre-molars (men)	chin
Incisors (men)	chin						
Canines (men)	chin						
Pre-molars (men)	chin						

AREATE or bald plates, is a common enough form of hair loss. Approximately 1% of the general population develop this problem at some time in their life. They are normally caused by nervous factors (stress, depressions, anxiety, etc.), and in a smaller degree by viral processes, mouth infections which can be infections of the mucous membrane or caries (see box below), and by thyroid problems which normally refer to a universal alopecia.



Treatment

Prepare the scalp by applying the Dermocapilar mask (ref. 28, 58) previously mixed with Herby (Ref. 25) and apply as if it was a dye, leave to act for 20 minutes, carry out this operation once a week on dirty hair.

Wash at least three times a week with Placent shampoo (Ref. 8) alternating with Royal Jelly shampoo (Ref. 35)
 Apply Wontersil Science 6 (Ref. 40) directly to the reflected areas twice a day for 90 days.